

# Motorized Spinal Decompression for Chronic Discogenic Low Back Pain: Chart Review of 100 Outpatients

As Presented at the:  
American Society of Anesthesiologists  
Oct. 15-17, 2006 | Chicago, IL



Alex Macario, MD, MBA, [Stanford University](#);  
Sunil J. Panchal, MD, [COPE Foundation](#), [Florida Pain Management](#);  
Charlotte Richmond, PhD, [Nema Research](#), [Biomedical Research & Education Foundation](#);  
Joseph V. Pergolizzi, Jr., MD, [Johns Hopkins University](#) & [Nema Research](#)

# Motorized Spinal Decompression for Chronic Discogenic Low Back Pain: Chart Review of 100 Outpatients

Alex Macario, MD, MBA, *Stanford University*;

Sunil J. Panchal, MD, *COPE Foundation, Florida Pain Management*;

Charlotte Richmond, PhD, *Nema Research, Biomedical Research & Education Foundation*

Joseph V. Pergolizzi, Jr., MD, *Johns Hopkins University & Nema Research*

## ABSTRACT

**Objective:** Conduct retrospective chart audit to assess outcomes of a random sample of outpatients treated with motorized spinal decompression via the DRX9000™ for chronic low back pain lasting more than 12 weeks.

**Methods:** Data from charts of 100 adults cared for in 2004-2006 at four clinics, one hospital-based and three free-standing, were abstracted using a standardized data collection form. Protected health information was accessed in accordance with the HIPAA privacy rule. Workman's compensation patients were excluded. DRX sessions (28-30 mins each) were for 8 weeks (mean) with 4-5 sessions the first week tapering to one session/wk (mean treatments = 23). Treatment protocol included instruction on lumbar stretching exercises and ice or muscle stimulation after DRX sessions. Pain, analgesic use, and activities of daily living were assessed pre and post treatment.

**Results:** Subjects (62% female, 94% white, mean age 55, 53% employed) had mean pain score 5.99 on a 0 to 10 scale (0=no pain 10=worst pain) at time of initial presentation that decreased to 0.87 after last DRX treatment. NSAID (41% of patients) and opioid (24% of the patients) use decreased (<5%) after treatment.

**Conclusion:** Overall, patients' pain improved after DRX treatment, requiring fewer analgesics, with better function. Practice variability exists in how clinics use the DRX9000™.

## OBJECTIVE

- Conduct retrospective chart audit to assess outcomes of a random sample of outpatients treated with motorized spinal decompression via the DRX9000™ for chronic low back pain lasting more than 12 weeks.

## METHODS

- Data from charts of 100 adults cared for in 2004-2006 at four clinics, one hospital-based and three free-standing, were abstracted using a standardized data collection form.
- Protected health information was accessed in accordance with the HIPAA privacy rule.
- Workman's compensation patients were excluded.
- DRX sessions (28-30 mins each) were for 8 weeks (mean) with 4-5 sessions the first week tapering to one session/wk (mean treatments = 23).
- Treatment protocol included instruction on lumbar stretching exercises and ice or muscle stimulation after DRX sessions.
- Pain, analgesic use, and activities of daily living were assessed pre and post treatment.

## RESULTS

- Subjects (62% female, 94% white, mean age 55, 53% employed) had mean pain score 5.99 on a 0 to 10 scale (0=no pain 10=worst pain) at time of initial presentation that decreased to 0.87 after last DRX treatment. NSAID (41% of patients) and opioid (24% of the patients) use decreased (<5%) after treatment (Fig. 1 - Fig 9).

**DEMOGRAPHICS**

Total Number of Patients = 100			
Mean Age	55	Mean Height	68 in
Female	62%	Mean Weight	89 kg
Employed	53%	White	94%
Retired	40%	Hispanic	3%
Disabled	5%	Black	2%
Housewife	1%	Asian	1%

**FIGURE 1**
**LIMITING EFFECTS OF LBP ON ACTIVITIES OF DAILY LIVING**

	Pre-DRX9000	Post-DRX9000
Bathing	25%	0%
Dressing	25%	0%
Walking	50%	1%
Sitting	50%	3%
Standing	51%	4%
Sleeping	21%	0%
None	1%	10%
Other	59%	1%
Unknown	9%	59%

**FIGURE 5**
**MEDICAL DIAGNOSIS & SYMPTOMS**

Medical Diagnosis		Symptoms	
Herniated Disc	74%	Nonspecific LBP	86%
Degenerative Disc Disease	66%	Leg Radiation	62%
Herniated & Degenerative Disc	26%	Radiation to Buttocks	22%
Sciatica	11%	Leg Pain > Back Pain	16%
Mean Duration Low Back Pain	260 wks	Prior Surgery	12%

**FIGURE 2**
**ANALGESIC USE**

Analgesic Use	Pre-DRX9000	Post-DRX9000
No meds	40%	20%
NSAIDs	43%	0%
Opioids	23%	0%
Muscle Relax	12%	1%
Steroids	4%	1%
Unknown	0%	59%

Cells do not add up to 100% as some patients were on more than one medication.

**FIGURE 6**
**MRI RESULTS PRE-TREATMENT**


MRI Results		Level of Pathology	
Disc bulge	36%	L5-S1	35%
Degenerative Changes	28%	L4-L5	40%
Protrusion	28%	L3-L4	13%
Extrusion	5%	L2-L3	8%
		L1-L2	1%
		T12-L1	2%

**FIGURE 3**
**POST DRX FOLLOW UP SCHEDULE**

0-3 months	3-6 months	6-9 months	> 9 months
50%	22%	9%	19%

**FIGURE 7**
**TREATMENT PROTOCOL AT SITES**

	Site A	Site B	Site C	Site D
Balance musculoskeletal system before DRX	Y	N	Y	Y
Heat before DRX	N	Y	N	N
Ice after DRX	Y	Y	Y	Y
Muscle stimulation after DRX	Y	N	Y	Y

**FIGURE 4**
**PAIN VRS**

**FIGURE 8**

**LIMITING EFFECTS OF LBP ON ACTIVITIES OF DAILY LIVING**

	Pre-DRX9000	Post-DRX9000
Bathing	25%	0%
Dressing	25%	0%
Walking	50%	1%
Sitting	50%	3%
Standing	51%	4%
Sleeping	21%	0%
None	1%	10%
Other	59%	1%
Unknown	9%	59%

FIGURE 5

**OSWESTRY LBP DISABILITY INDEX**




FIGURE 9

**ANALGESIC USE**

Analgesic Use	Pre-DRX9000	Post-DRX9000
No meds	40%	20%
NSAIDs	43%	0%
Opioids	23%	0%
Muscle Relax	12%	1%
Steroids	4%	1%
Unknown	0%	59%

Cells do not add up to 100% as some patients were on more than one medication.

FIGURE 6

**RESULTS CONTINUED**

- Patients reported a mean 90% improvement in back pain, and better function as measured by activities of daily living. On a 0 to 10 scale (0=Not satisfied 10=Very satisfied) patients rated the DRX9000™ an 8.98 (Fig. 10).
- No patient required more invasive therapies (e.g., surgery).

**POST DRX FOLLOW UP SCHEDULE**

	0-3 months	3-6 months	6-9 months	> 9 months
	50%	22%	9%	19%

FIGURE 7

**PATIENT SATISFACTION**

Mean satisfaction with DRX (0-10 scale) 0=not satisfied 10=Very satisfied	8.98
Improvement in LBP provided by DRX	90%
Recommend DRX to someone else	100%

Data based on 20-25% of patients contacted in follow up

FIGURE 10

**PAIN VRS**




FIGURE 8

**CONCLUSION**

Overall, patients' pain improved after DRX treatment, requiring fewer analgesics, with better function. Practice variability exists in how clinics use the DRX9000™. We didn't have control groups, making it difficult to know how much of the benefit was placebo or spontaneous recovery and how much was due to the intervention. Randomized double-blinded clinical trials are needed to measure the efficacy of non-surgical spinal decompression systems.